



# Prediction of Personal Glycemic Responses to Food for Individuals With Type 1 Diabetes Through Integration of Clinical and Microbial Data

**Document Version:** Accepted author manuscript (peer-reviewed)

# Citation for published version:

Shilo, S, Godneva, A, Rachmiel, M, Korem, T, Kolobkov, D, Karady, T, Bar, N, Wolf, BC, Glantz-Gashai, Y, Cohen, M, Zuckerman Levin, N, Shehadeh, N, Gruber, N, Levran, N, Koren, S, Weinberger, A, Pinhas-Hamiel, O & Segal, E 2022, 'Prediction of Personal Glycemic Responses to Food for Individuals With Type 1 Diabetes Through Integration of Clinical and Microbial Data', *Diabetes Care*, vol. 45, no. 3, pp. 502-511. https://doi.org/10.2337/dc21-1048

*Total number of authors:* 18

Digital Object Identifier (DOI): 10.2337/dc21-1048

Published In: Diabetes Care

License: CC BY

# **General rights**

@ 2020 This manuscript version is made available under the above license via The Weizmann Institute of Science Open Access Collection is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognize and abide by the legal requirements associated with these rights.

How does open access to this work benefit you?

Let us know @ library@weizmann.ac.il

# Take down policy

The Weizmann Institute of Science has made every reasonable effort to ensure that Weizmann Institute of Science content complies with copyright restrictions. If you believe that the public display of this file breaches copyright please contact library@weizmann.ac.il providing details, and we will remove access to the work immediately and investigate your claim.



https://doi.org/10.2337/dc21-1048



Smadar Shilo,<sup>1,2,3</sup> Anastasia Godneva,<sup>1,2</sup> Marianna Rachmiel,<sup>4,5</sup> Tal Korem,<sup>1,2,6</sup> Dmitry Kolobkov,<sup>1,2</sup> Tal Karady,<sup>1,2</sup> Noam Bar,<sup>1,2</sup> Bat Chen Wolf,<sup>1,2</sup> Yitav Glantz-Gashai,<sup>3</sup> Michal Cohen,<sup>3,7</sup> Nehama Zuckerman Levin,<sup>3,7</sup> Naim Shehadeh,<sup>3,7</sup> Noah Gruber,<sup>5,8</sup> Neriya Levran,<sup>8,9</sup> Shlomit Koren,<sup>5,10</sup> Adina Weinberger,<sup>1,2</sup> Orit Pinhas-Hamiel,<sup>5,8</sup> and Eran Segal<sup>1,2</sup>

## OBJECTIVE

Despite technological advances, results from various clinical trials have repeatedly shown that many individuals with type 1 diabetes (T1D) do not achieve their glycemic goals. One of the major challenges in disease management is the administration of an accurate amount of insulin for each meal that will match the expected postprandial glycemic response (PPGR). The objective of this study was to develop a prediction model for PPGR in individuals with T1D.

# **RESEARCH DESIGN AND METHODS**

We recruited individuals with T1D who were using continuous glucose monitoring and continuous subcutaneous insulin infusion devices simultaneously to a prospective cohort and profiled them for 2 weeks. Participants were asked to report real-time dietary intake using a designated mobile app. We measured their PPGRs and devised machine learning algorithms for PPGR prediction, which integrate glucose measurements, insulin dosages, dietary habits, blood parameters, anthropometrics, exercise, and gut microbiota. Data of the PPGR of 900 healthy individuals to 41,371 meals were also integrated into the model. The performance of the models was evaluated with 10-fold cross validation.

#### RESULTS

A total of 121 individuals with T1D, 75 adults and 46 children, were included in the study. PPGR to 6,377 meals was measured. Our PPGR prediction model substantially outperforms a baseline model with emulation of standard of care (correlation of R = 0.59 compared with R = 0.40 for predicted and observed PPGR respectively;  $P < 10^{-10}$ ). The model was robust across different subpopulations. Feature attribution analysis revealed that glucose levels at meal initiation, glucose trend 30 min prior to meal, meal carbohydrate content, and meal's carbohydrate-to-fat ratio were the most influential features for the model.

# CONCLUSIONS

Our model enables a more accurate prediction of PPGR and therefore may allow a better adjustment of the required insulin dosage for meals. It can be further implemented in closed loop systems and may lead to rationally designed nutritional interventions personally tailored for individuals with T1D on the basis of meals with expected low glycemic response. <sup>1</sup>Department of Computer Science and Applied Mathematics, Weizmann Institute of Science, Rehovot. Israel

<sup>2</sup>Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel

<sup>3</sup>Pediatric Diabetes Clinic, Institute of Diabetes, Endocrinology and Metabolism, Rambam Health Care Campus, Haifa, Israel

<sup>4</sup>Pediatric Endocrinology Unit, Shamir Medical Center, Zerifin, Israel

<sup>5</sup>Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel

<sup>6</sup>Department of Systems Biology, Columbia University, NY

<sup>7</sup>Bruce Rappaport Faculty of Medicine, Technion, Haifa, Israel

<sup>8</sup>Pediatric Endocrine and Diabetes Unit, Edmond and Lily Safra Children's Hospital, Sheba Medical Center, Ramat-Gan, Israel

<sup>9</sup>Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot, Israel

<sup>10</sup>Diabetes Unit, Shamir Medical Center, Zerifin, Israel

<sup>a</sup>Corresponding authors: Eran Segal, eran. segal@weizmann.ac.il, and <sup>b</sup>Orit Pinhas-Hamiel, orit.hamiel@sheba.health.gov.il

Received 15 May 2021 and accepted 17 September 2021

Clinical trial reg. no. NCT02919839, clinicaltrials.gov

This article contains supplementary material online at https://doi.org/10.2337/figshare.16649266.

© 2021 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at https://www. diabetesjournals.org/content/license.

Diabetes Care Publish Ahead of Print, published online October 28, 2021

Type 1 diabetes (T1D) is one of the most common chronic diseases in children. Despite technological advances introduced in the past decades, such as continuous glucose monitoring (CGM) and continuous subcutaneous insulin infusion (CSII) devices (1), results from various large-scale clinical trials repeatedly show that the clinical management of T1D is challenging, especially in children and adolescents, with many patients not achieving the glycemic goals recommended by clinical guidelines (2-4). One of the greatest challenges in disease management is the administration of an accurate amount of insulin for each meal. While high levels of insulin might result in a life-threatening hypoglycemic event, a low level would result in high postprandial blood glucose levels and its associated short-term and long-term comorbidities (5). The Standards of Medical Care in Diabetes guidelines, published by the American Diabetes Association, state that patients should match prandial insulin to carbohydrate intake, premeal blood glucose, and anticipated physical activity (6). However, it was previously shown that conventional therapy resulted in suboptimal insulin counteraction of postprandial glycemic responses (PPGRs) (7). More sophisticated models exist, such as those that also include adjustment for the meal's fat and protein content, but have so far failed to provide a significant improvement in glycemic control, and their implementation in real practice is limited (8,9).

A previous study by Zeevi et al. (10) in healthy individuals revealed that carbohydrate content alone is not a good predictor of glycemic responses to meals and that a significant interindividual variability in PPGR exists. Moreover, environmental factors, including the gut microbiota, were associated with the glycemic response of healthy individuals to meals (10). We therefore hypothesized that by collecting data on the administered insulin dosages, along with additional clinical and microbial data that were previously shown to contribute to PPGR prediction in healthy individuals, we will be able to construct a prediction model for glycemic responses to meals specifically tailored for individuals with T1D. An accurate model may have an immense clinical value, as it will allow a better estimation of the required insulin dosage for meals and can guide personal nutritional interventions, which will be

based on meals with expected low glycemic response.

# RESEARCH DESIGN AND METHODS Study Design

To characterize the PPGRs of individuals with T1D, we conducted a prospective observational study. At study initiation, a physician authorized participation and acquired informed consent. Anthropometric measurements and vital signs were taken by the medical staff, and blood tests, including metabolic and lipid profile, HbA<sub>1c</sub>, and thyroid function, were drawn and analyzed in the laboratories of each of the three medical institutes. Health and lifestyle questionnaires were filled out by the participants or their legal guardians (Supplementary Fig. 1). During the 2 weeks of study participation, participants continued using their own personal CGM and CSII devices. Participants who did not have a CGM device in their possession were connected to a Free-Style Libre Flash CGM (FSL-CGM) system for the duration of the study. Continuous glycemic and insulin profiles were then acquired in high resolution from the insulin pump and CGM records, with 5- to 15-min intervals between glucose measurements, depending on the CGM type.

Study participants, or their parents in the case of young children, used a proprietary smartphone app (www.personal nutrition.org) (Supplementary Fig. 2), already used by >900 people (10), to log, in real time, food intake, sleep times, physical activity, and medication intake with the exception of insulin, which was recorded in the CSII devices. Each food item within every meal was logged, along with its weight by selecting it from a database of 6,401 foods with full nutritional values based on the Israeli Ministry of Health database with the expansion of additional items by certified dietitians. For increased compliance, participants were informed that accurate logging is crucial for them to receive an accurate analysis of their PPGRs to food. Participants were asked to follow their normal daily routine and dietary habits, with the exception of seven standardized meals.

# **Participant Recruitment**

Enrollment and recruitment of participants were conducted by the medical teams in three medical centers in Israel: Sheba Medical Center, the largest hospital in Israel; Rambam, Health Care Campus, Northern Israel's primary hospital; and Shamir Medical Center, the fourthlargest government hospital in Israel. The locations of these clinics allowed us a nearly national coverage of the Israeli population. The inclusion criteria were age between 3 and 70 years, >1 year since T1D diagnosis, use of CGM and CSII devices simultaneously, and a capability to work with a mobile phone app on a daily basis for the recording of the dietary intake by participants or their parents in the case of young children. A minimal age of 3 years was chosen, since it was previously shown that in this age the development of the microbiome from infancy reaches a stable phase (11). A maximal age of 70 years was chosen due to the higher prevalence of additional comorbidities in this age-group. Participants were recruited at least 1 year following diagnosis for minimization of the possible effects of the "honeymoon period" of T1D in which a good glycemic control is achieved with reduced insulin requirements (9). Exclusion criteria included an active inflammatory or neoplastic disease, pregnancy, and antibiotic usage 3 months prior to participation in the study.

#### Meals Preprocessing

We applied the following consecutive filters on the 9,139 meals that were logged by the participants: First, we merged meals logged <30 min apart if the earliest meal contained >50 cal. The merged meal was assigned the summed values of all of its components and the time of the earliest component. Second, we removed 1) meals logged within 90 min of other meals to avoid the potential influence of adjacent meals and their preceding insulin administration. 2) meals with components weighing >1 kg, 3) meals with incomplete logging, 4) meals with <70 kcal and <15 g total weight, and 5) meals with carbohydrate content >200 g, baseline insulin level of 0, or baseline glucose level <50 mg/dL (2.8 mmol/L). Glucose level at baseline was considered as the lowest glucose value within ±15 min from self-reporting of the meal in the app. This process resulted in 6,377 meals that were logged during the study and were included in the analysis.

#### **Standardized Meals**

Each subject received seven standardized breakfasts, provided by the study team, to consume throughout the 14 days of the study:  $2 \times 30$  g glucose,  $2 \times$ 50 g glucose, 2  $\times$  65 g bread, and 65 g bread + 20 g butter. Participants were instructed to consume these meals immediately after their night fast, not to modify the meal, and to refrain from eating or performing strenuous physical activity before and for 2 h following meal consumption. In addition, they were instructed to administer insulin calculated by their personal carbohydrate-to-insulin ratio, insulin sensitivity factor, and glycemic target. Participant adherence for consuming the standardized meals was partial, with 116 people consuming at least one test meal, 48 consuming 2 × 30 g glucose, 51 consuming 2 × 50 g glucose, 35 consuming  $2 \times 65$  g bread, and only 3 consuming 65 g bread + 20 g butter twice. Overall, 688 standardized meals were consumed during the study. Following meal preprocessing described above, 118 meals were included in the analysis. Since insulin boluses are calculated by the carbohydrate content of the meal and by the baseline glucose level (12), we compared the within-individual glycemic response of two identical standardized meals only when the difference between glucose levels at meal initiation was <40 mg/dL (2.2 mmol/L). In addition, we only included meals consumed in the morning (5:00-10:00 A.M.).

# Prediction of Prandial Glucose Response

To measure the glucose response to meals we used two matrices. First, the PPGR of each meal was calculated by combination of the reported meal time with CGM data and computation of the incremental area under the glucose curve in the 2 h after the meal (13). Of note, no significant differences between PPGRs extracted from CGMs and those obtained from blood tests were shown in healthy individuals in a previous study (14). Second, we calculated the difference between glucose level at meal initiation and maximal glucose level during the 2 h after the meal (Glumax) for the multiple real-life meals. This measure

was chosen because it is less sensitive to inaccurate logging time by the participants. For predicting both indices, we constructed prediction models based on XGBoost 0.90; 3,000 estimators and a learning rate of 0.005 were used.

A total of 296 features were included in the model as input, including features representing meal content and blood tests results, CGM and insulin pumpderived features, questionnaires, and microbiome features (Supplementary Table 1). To test the performance of our models, we evaluated our model using several 10-fold cross validation schemes. The results presented in the manuscript are from a per-meal validation scheme, in which the model was trained on 90% of the meals and the remaining 10% of the meals were used as validation. Additional cross-validation schemes were analyzed (Supplementary Fig. 4). Pearson R between predicted and observed PPGR and Glu<sub>max</sub> was calculated for each model. Explained variance was computed as  $R^2$  (coefficient of determination) regression score function in the Python scikit-learn library.

## Integration of Data From Healthy Individuals

To analyze whether an integration of data of postprandial responses to meals will improve the prediction results, we integrated data of the PPGR values of 41,371 meals from 900 healthy individuals. The cohort characteristics are described in detail in the study by Zeevi et al. (10) from 2015. In brief, the healthy cohort included individuals age 18–70 years, not previously diagnosed with diabetes, who logged meals in real time in the same smartphone app used in the current study and were connected to a CGM device for 1 week. The clinical data collected from participants were similar with the exception of insulin dosage, which is not relevant for healthy individuals. We used the data from the healthy cohort in two schemes. First, we tested the performance of a model trained solely on data from the healthy individuals on the cohort of individuals with T1D. Second, we tried using the output of a prediction model trained on the data that originates from healthy individuals as an additional feature to our model, based on data of individuals with T1D. We evaluated the value of Pearson correlation obtained between

predicted and measured PPGR following these two approaches.

#### **Feature Attributions**

We used the recently introduced SHapley Additive exPlanation (SHAP) methods (15–17) for model interpretability. SHAP values are calculated individually for every feature and represent the average change in the model's output upon conditioning on that feature, when introducing each feature separately, as it is introduced one at a time. The additive property of SHAP values was used to analyze the impact of different groups of features on the model.

#### Stool Sample Collection and Genomic DNA Extraction

Participants were instructed to sample their stool once during the study period, following detailed printed instructions. Sampling was done using both a swab and an OMNIgene GUT (OMR-200; DNA Genotek) stool collection kit. Collected samples were immediately stored in a home freezer ( $-20^{\circ}$ C), and transferred in a provided cooler to our facilities where they were stored at  $-80^{\circ}$ C ( $-20^{\circ}$ C for OMNIgene GUT kits) until DNA extraction. Gut microbiota profile was obtained only from the DNA Genotek samples by metagenomics sequencing.

Metagenomic DNA was purified with the DNeasy PowerMag Soil DNA extraction kit (QIAGEN), optimized for Tecan automated platform. Next-generation sequencing libraries were prepared with use of Nextera DNA Library Prep (Illumina) and sequenced on a NovaSeq sequencing platform (Illumina). Sequencing was performed with 100-base pairs single-end reads with depth of 10 million reads per sample. We filtered metagenomic reads containing Illumina adapters, filtered low-quality reads, and trimmed low-quality read edges. We detected host DNA by mapping with Bowtie2 (18) to the human genome with inclusive parameters and removed those reads. Bacterial relative abundance (RA) estimation was performed through mapping bacterial reads to species-level genome bins of representative genomes (19). We selected all species-level genome bins representatives with at least five genomes in a group and for these representatives genomes kept only unique regions as a reference data set. Mapping was performed with Bowtie2 (18), and we estimated abundance by calculating the mean coverage of unique genomic regions across the 50% most densely covered areas as previously described (20). Feature names include the lowest taxonomy level identified.

Δ

62 Sheba medical center

antibiotic usage

1 Excluded due to a recent

#### **Ethics Approval**

The study was approved by the Rambam Health Care Campus institutional review board (IRB), Tel Hashomer Hospital IRB, Shamir Medical Center IRB, and Weizmann Institute of Science IRB. All participants signed written informed consent forms. All identifying details of the participants were removed prior to the computational analysis. The trial was registered at ClinicalTrials.gov, clinical trial reg. no. NCT02919839.

#### Data and Resource Availability

Metagenomic sequencing data that support the findings of this study will be made available. Clinical data cannot be shared due to restrictions by virtue of the informed consent. The data set is available from https://data.mendeley.com/data sets/bcz47mhvc3/1. Analysis code is available from https://github.com/nastyagod neva/t1d\_microbiome.

#### RESULTS

#### Study Population

A total of 142 individuals with T1D were recruited into the study between March 2017 and April 2019. Five individuals (3.5%) dropped out, resulting in 137 participants, of whom 131 provided data from CGM and CSII devices. Of these, 121 individuals, 46 children (<18 years of age) and 75 adults, logged meals during the study period and were included in the analysis (Fig. 1A). The mean ± SD age was 24.7 ± 15 years (median 21 years, interquartile range [IQR] 14-32) (Fig. 2A), and average disease duration was 11.2 ± 10.1 years (median 8 years IQR 4–15). Mean HbA<sub>1c</sub> level was 7.5% ± 1.1% (58.5 ± 12.1 mmol/mol) (Fig. 2B; see Supplementary Table 2 for the results of all blood tests obtained at baseline). Mean BMI of adults was 25.1 ± 3.9 kg/m<sup>2</sup> (median 22.4 kg/m<sup>2</sup> IQR [20.4–22.6]), and mean BMI percentile of children when converted according to the Centers for Disease Control and Prevention reference percentiles was 58 ± 23 (median 59.4 IQR[38.3–83.5]) (Supplementary Fig. 3). Of the 121 participants, 42 (34.7%) had



Recruitment to the study

March 2017 to April 2019

47 Rambam medical center

1 Excluded due to a recent

antibiotic usage

36 Shamir medical center

1 Excluded due to a recent

antibiotic usage

**Figure 1**—A: Cohort selection; number of participants with T1D is indicated. *B*: Study scheme. Data integrated into the prediction model for the glycemic response to meals are presented. The prediction model was constructed for 1) PPGR and 2) Glu<sub>max</sub> in individuals with T1D. All data types were available for all participants, except for stool samples, which were included for 106 individuals, and physical activity which was logged by 83 participants. HbA<sub>1c</sub> value was missing in one blood test. AUC, area under the curve.

at least one additional comorbidity. The most common comorbidities were hypothyroidism (17 participants [14%]), hyperlipidemia (13 participants [10.7%]), and celiac disease (7 participants [5.8%]). Thirty-three participants (27%) consumed additional medications apart from insulin during the study. The most common medications were levothyroxine, oral contraceptives, and antilipidemic drugs. (See Supplementary Table 3 for a full list of medical conditions and medications.) None of the participants reported using long-acting insulin during the study. Altogether, 1,775 days of a concurrent usage of both CGM (including 302,703 glucose measurements) and CSII devices simultaneously were included in the analysis. Of them, 1,597 days included at least one meal logging. A total of 110 individuals provided a stool sample including 7 participants who had celiac disease and were excluded from all the microbiome

analyses. Cohort characteristics are presented in Table 1.

#### **Nutritional Profiling**

Overall 9,139 meals (mean of 73.9 meals per participant, median 68 meals, [IQR 51.75-93), a total of 2,383,181 kcal (mean of 19,695 kcal per participant, median 18,631 kcal, IQR 14,517-24,903), were logged by participants during the study. The distributions of the number of meals logged for children and adults are presented in Supplementary Fig. 3. To obtain a global view of the dietary habits of the participants, we first analyzed the fraction that each food group contributed to the cohort's overall energy intake (Fig. 2D and E) and the distribution of macronutrients intake from the overall energy intake (Fig. 2C). The main food groups consumed by the participants were similar in children and adults. Average carbohydrate, fat, and protein consumption



**Figure 2**—Clinical and nutritional data. In *A*–*C*, adults are marked in orange and children in blue. Distribution of age (*A*) and HbA<sub>1c</sub> (%) measured in a blood test (*B*) at baseline. *C*: Distribution of the average daily macronutrient composition from the total energy intake for individuals in the cohort (each participant is represented by a dot). *D* and *E*: Major food components consumed by energy intake are presented for adults (*D*) and children (*E*).

was 39.1 ± 8%, 41.6 ± 10.2%, and 17.1 ± 4.3% of total energy, respectively (Table 1 and Fig. 2). Overall, macronutrients distribution changed with age, with older individuals consuming fewer carbohydrates, as demonstrated by a positive association between age and the percentage of lipids (r = 0.27, P < 0.05, false discovery rate [FDR[ corrected) and negative association of age with the percentage of carbohydrates from total energy intake (r =-0.30, P < 0.05, FDR corrected). After adjustment for age, the percentage of carbohydrates from the total energy intake consumed per day by the participants was positively correlated with blood levels of HbA<sub>1c</sub> at study initiation (r =0.27, P < 0.05, FDR corrected), mean glucose value (r = 0.21, P < 0.05, FDR corrected), and glucose variability (r = 0.27, P < 0.05, FDR corrected) (Supplementary Fig. 4).

# Glycemic Responses of the Same Person to the Same Meal Are Reproducible

We next analyzed intrapersonal variability in the PPGR of the same person to the same meal. To that end, we assessed whether the PPGRs of standardized meals that were given twice to each participant are reproducible. Overall, 688 standardized meals were consumed during the study by 116 participants. However, to limit cases in which an independent exposure leads to the variability, we performed this analysis only for standardized meals consumed in the morning, in which case the baseline glucose levels at meal initiation were similar (see Research Design and Methods). Altogether, 118 test meals were included for 59 individuals. In these cases, there was a relatively high agreement for all duplicated test meals (R = 0.63), demonstrating that the PPGR to identical meals is correlated in the same participant when the baseline glucose level is similar (Supplementary Fig. 5).

# Prediction of Glycemic Responses to Real-life Meals

We next examined our ability to predict 1) PPGR and 2)  $Glu_{max}$ , which we defined as the maximal glucose rise during the 2 h after meal initiation

(Fig. 1B). After meal preprocessing, from 9,139 logged meals, 6,377 meals (mean ± SD 47.4 ± 24.1 meals per participant [Supplementary Fig. 3]) were included in the analysis. (See Research Design and Methods). For both tasks, we constructed prediction models based on extreme gradient boosting (XGBoost) taking as input the following: 1) Only the meal's carbohydrate content. 2) A baseline model representing standard of care for insulin administration. Since insulin boluses are calculated on the basis of the carbohydrate content of the meal and the baseline glucose level, the meal's carbohydrate content and glucose at meal initiation were included in the model (12). Insulin bolus given 90 min prior to the meal was also included in the baseline model, for cases in which it was already administered for the meal, thereby influencing the glycemic response. And 3) all the information that was collected from the participant during the study period, with regard to features representing meal content (e.g., macronutrients), meal timing, daily activity (e.g., time from logged exercise), blood

|  | All (n = 121)   | Adults ( <i>n</i> = 75)  | Children ( $n = 46$ )                                 |
|--|---|--|---|
| Age (years)  | 24.7 ± 15   | 32.3 ± 14.1  | 12.1 ± 3.7  |
| Diabetes duration (years)  | 11.2 ± 10.1   | 15.1 ± 10.7  | 4.5 ± 2.7   |
| Male sex   | 40  | 40   | 40  |
| Weight (kg)  | 61.9 ± 19.5   | 71.4 ± 12.9  | 46.1 ± 18.3   |
| BMI (kg/m <sup>2</sup> )   | 23.1 ± 4.8  | 25.1 ± 3.9   | 19.8 ± 4.2  |
| BMI percentile, for children   |   |  | 58 ± 23   |
| HbA <sub>1c</sub> (%, mmol/mol)  | 7.5 ± 1.1, 58.5 ± 12.1                                      | 7.3 ± 1.0, 56.3 ± 10.9   | 7.8 ± 1.3, 61.7 ± 14.25                               |
| Microbiome samples (N)   | 110   | 71   | 39  |
| Individuals who logged physical activity (N)   | 83  | 58   | 25  |
| Real-time meal logging (per participant)<br>Days of meals logging + CGM + CSII data<br>Meals logged<br>Meals included in the predictor<br>Total energy intake per day (kcal) | 13.2 ± 3.3<br>73.9 ± 32.2<br>52.7 ± 24.0<br>1,516.7 ± 550.2 | $13.2 \pm 3.1 \\76.9 \pm 27.3 \\55.3 \pm 25.6 \\1,571.6 \pm 523.4$ | 13.4 ± 3.4<br>69 ± 26.2<br>48.5 ± 21<br>1,428.4 ± 585 |
| Nutrition components<br>% total energy intake from lipids<br>% total energy intake from protein<br>% total energy intake from carbohydrates                                  | 39.1 ± 8<br>17.1 ± 4.3<br>41.6 ± 10.2                       | 40.3 ± 7.6<br>17.5 ± 4.8<br>39.6 ± 9.8                             | 37.1 ± 8.5<br>16.2 ± 3.3<br>45.2 ± 10                 |

# Table 1–Cohort characteristics

Data are means  $\pm$  SD or % unless otherwise indicated. Characteristics of individuals with T1D who were included in the prediction model are presented. BMI percentiles for children were converted to reference percentiles provided by the Centers for Disease Control and Prevention.

parameters (e.g.,  $HbA_{1c}$ ), CGM-derived features, data on insulin dosage obtained from CSII devices, questionnaires, and microbiome features (e.g., metagenomic RA). (A full list of features can be found in Supplementary Table 1.) Several cross-validation schemes were used for validation of the model (see *Research Design and Methods* and Supplementary Fig. 6).

For the prediction of PPGR, a model that is based solely on the meal's carbohydrate content achieves a relatively low correlation with PPGRs (R = 0.16) and explains only  $\sim$ 3% of the variance in glycemic response (Fig. 3A). A baseline model using carbohydrate content, glucose level at meal initiation, and insulin bolus performs better and explains  $\sim$ 16% of the variance in glycemic response (R = 0.4,  $P < 10^{-10}$ ) (Fig. 3B). The full model, with integration of glucose measurements, insulin dosages, dietary habits, blood parameters, anthropometrics, exercise, and gut microbiota (Supplementary Table 1), achieves a substantially higher correlation for the held out PPGRs of meals and increases the explained variance to  $\sim$ 35% (R = 0.59,  $P < 10^{-10}$ ) (Fig. 3C). Similarly, for Glu<sub>max</sub> prediction, a meal's carbohydrate-based model achieves a relatively low correlation

(R = 17) (Fig. 3*D*), a baseline model performs better (R = 0.43,  $P < 10^{-10}$ ) (Fig. 3*E*), and the full model predicts the held out values of individuals with a significantly higher correlation (R = 0.61,  $P < 10^{-10}$ ) (Fig. 3*F*).

Although insulin doses substantially affect glucose levels in individuals with T1D, we next asked whether integration of data on glycemic responses to meals from healthy individuals may also contribute to our ability to predict the glycemic response to meals in T1D. To this end, we used an extensive data set of detailed clinical profiling and PPGR measurements of 41,371 meals from 900 healthy individuals (10) (see Research Design and Methods). First, we tested a model trained solely on data from the healthy individuals cohort on the T1D cohort. This model achieved a correlation between predicted and measured PPGR of R = 0.39, similar to that achieved by our baseline model but lower than achieved by the full model trained specifically on individuals with T1D. Second, we tried using the output of a prediction model trained on the data that originate from healthy individuals as an additional feature to our model, which is based on data of individuals with T1D. This resulted in a correlation similar to that obtained by the original model (Supplementary Fig. 7).

## Variability in the Prediction of Glycemic Responses

To further investigate the performance of the full model, we analyzed the variability in the correlation between predicted and observed PPGR between individuals in the cohort. While in most of the individuals, a correlation of >0.52 was obtained, a high variability was observed (Supplementary Fig. 3). To further analyze the factors underlying this variability and the overall robustness of the model, we divided the cohort into several subgroups by clinical parameters and calculated the correlation for each subgroup. Subgroup analysis revealed similar results for divisions by age, HbA1c, time spent in a state of hypoglycemia, and different types of CGM and CSII devices (Supplementary Table 4). We next compared the characteristics of the participants with a high correlation with the characteristics of those with a low correlation and found that the latter had a significantly higher glucose variability and logged significantly fewer meals per day (3.58 vs. 4.26, P <0.05, FDR corrected) and throughout the



**Figure 3**—PPGR and Glu<sub>max</sub> predictions for multiple real-life meals. Dots represent predicted (*x*-axis) and CGM-measured 1) PPGR (*y*-axis) for a model based on only the meal's carbohydrate content (*A*), the meal's carbohydrate content, glucose level at meal initiation, and bolus insulin administered 90 min prior to the meal (*B*), and all features (*C*) and 2) Glu<sub>max</sub> (*y*-axis) for a model based on only the meal's carbohydrate content (*D*), the meal's carbohydrate content, glucose level at meal initiation, and bolus insulin administered 90 min prior to the meal (*B*), and all features (*C*) and 2) Glu<sub>max</sub> (*y*-axis) for a model based on only the meal's carbohydrate content (*D*), the meal's carbohydrate content, glucose level at meal initiation, and bolus insulin administered during the 90 min prior to the meal (*E*), and all features (*F*). All models are evaluated in 10-fold cross validation on the entire T1D cohort; explained variance values are indicated. *G*: Interpretation of the prediction model. SHAP values (in absolute log-odds scale) of the 10 most impactful features on the PPGR prediction model. Numbers near features name represent the minutes prior to meal initiation and glucose level 30 min prior to the meal. *H*: SHAP values (in absolute log-odds) of the 10 most impactful groups of features are presented as follows: "CGM" and "Insulin" groups include glucose values obtained from CGM devices and insulin dosages recorded by the insulin pump, respectively; "Nutrients" includes the nutritional content of the meal; "Microbiome" includes RA of bacteria detected from stool samples; and "Routine" includes the time interval from the latest physical activity reported and time of day in which the meal was consumed. carbs, carbohydrates.

study period (54.68 vs. 42.81, P < 0.05, FDR corrected) (Supplementary Table 5).

# Factors Underlying the Prediction of Postprandial Glycemic Responses

To gain insight into factors affecting prediction, we performed feature attribution analysis using SHAP (15–17). SHAP values represent the average change in the model's output upon conditioning on a specific feature (see *Research Design and Methods*). The most influential features with the highest mean absolute SHAP value included glucose levels at meal initiation, glucose trend in the 30 min prior to meal, meal carbohydrate content, the ratio between carbohydrate and fat in the meal, and time of day, for which the impact of earlier hours in the day was toward a higher glycemic response (Fig. 3*G*). By adding up the SHAP values

**Diabetes Care** 

of related features, we could analyze the overall impact of groups of features to the model. These analyses revealed that glucose measurements were the most impactful in the prediction, followed by the meal composition, insulin dosage, and microbial composition (Fig. 3*H*).

# CONCLUSIONS

In this study we used comprehensive clinical data from a 2-week profiling period to devise a prediction model for glycemic responses to meals in individuals with T1D. PPGR is an important contributor to the overall glycemic control (21), and decreasing it may improve the time spent in the target glycemic range, which is strongly associated with the risk of future microvascular complications in individuals with T1D (22). In recent years, the increased availability of sequential glucose measurements and insulin data, as a result of an increased usage of CGM and CSII devices, respectively, has paved the way for machine learning approaches targeted at the prediction of the dynamics of blood glucose in T1D (23). However, modeling the effect of different meals remains a challenge (23,24). To the best of our knowledge, our study is the first to investigate the contribution of adding data on PPGRs originating from healthy individuals and to incorporate microbial features to a prediction model for PPGR of individuals with T1D.

Here, we have shown a relatively high agreement (R = 0.63) for duplicated test meals of the same individual with T1D, demonstrating that the PPGR to identical meals is correlated within the same person when baseline glucose level is similar (Supplementary Fig. 5). However, it is worthy of note that the same analysis previously performed on a larger cohort of healthy individuals achieved substantially better results (R = 0.77 for glucose and bread with butter test meals, R = 0.71 for bread) (10). This emphasizes the fact that prediction of glycemic responses to meals is more challenging in individuals with T1D than in healthy individuals, as those with T1D have a higher glucose variability; are treated with insulin, which can greatly impact glucose levels; and they may engage in physical activity in proximity to the meal, which may also greatly impact their glucose levels.

Our prediction model for the glycemic response to real-life meals takes into account a comprehensive clinical and microbiome profile as input and aims at predicting two glycemic measures: PPGR and Glumax. In both glycemic measures, our model substantially outperforms a baseline model that emulates the current Standards of Care for insulin administration for meals in individuals with T1D and takes into account only the carbohydrate content of the meal and glucose level at meal initiation (6), as well as the dosage of bolus insulin already given prior to the meal (Fig. 4). Notably, a model based solely on the meal's carbohydrate content achieved a relatively low correlation with PPGRs (R = 0.16) compared with the correlation previously reported for healthy individuals (R = 0.38) (10), possibly due to the fact that the insulin dosage given for the meal has already been calculated and adjusted for the carbohydrate content. Integration of data from 900 healthy individuals (10) into the model did not improve its performance.

While a high variability in the correlation of predicted and observed glycemic response to meals was observed between participants, subgroup analysis has shown that the model is overall robust and performs similarly across individuals in different age-groups and with different levels of glycemic control. Glucose variability was significantly lower in individuals who had a high correlation, reflecting the challenge of glucose prediction in patients with high variability in glucose values. Notably, the number of meals logged per day and throughout the study was significantly higher in individuals who had a high correlation. This may be due to better adherence to the study protocol, which resulted in a more accurate and timely meal logging that enabled a better prediction. In addition, the fact that the model was trained on a larger number of meals logged by these participants may have also improved its performance. Nonetheless, these findings emphasize the potential to achieve a better prediction with additional accurate and timely meal logging in individuals with insufficient data.

Analysis of the impact of different factors on the prediction model revealed insights on the drivers of glucose rise following meals in individuals with T1D. The most impactful features included

glucose at meal baseline and the carbohydrate content of the meal, which are currently the main factors taken into consideration in the calculation of insulin bolus (6), thus further providing validation to our analyses. Other impactful factors, which are not taken into account in the Standards of Care for insulin administration today, are the glucose trend 30 min prior to the meal start and the ratio of carbohydrate to fat in the meal. The impact of earlier hours in the day was toward a higher glycemic response, which is in line with previous studies on diurnal changes in insulin requirements that showed that more insulin is needed per carbohydrate count (a lower carbohydrate-to-insulin ratio) in breakfast versus lunch and dinner (25,26). Overall, glucose values prior to the meal, followed by meal composition and insulin dosages, were the most impactful groups of features in the model. Interestingly, analyses of the impact of groups of features on the model revealed that glucose measurements were the most impactful in the prediction, followed by the meal composition, insulin dosage, and microbial composition (Fig. 3H).

In this cohort, individuals who consumed a diet with fewer carbohydrates had better glycemic control, as demonstrated by a higher percentage of time in range and a lower glucose average. The percentage of carbohydrates from the total energy intake consumed by the participants was positively correlated with blood levels of HbA<sub>1c</sub> at study initiation and with a higher glucose variability. Although several studies reported a statistically significant reductions in HbA<sub>1c</sub> in individuals with T1D consuming low-carbohydrate diets, others did not, and the overall effect of these diets is still debatable, partially due to heterogeneity of studies in this field (27).

This study has several limitations. First, a low adherence for standardized meals consumption and inaccurate selfreports of meals by participants could have affected our ability to predict the glycemic response to meals, and we believe that better predictions can likely be achieved by training the model on a larger volume of high-quality clinical data. This is empirically shown by the superiority of the model performance in participants who logged more meals and were more adherent to the study protocol (Supplementary Table 2). This limitation also presents a challenge in real-life utility of the model, as an accurate input of meals is required. Second, we did not have information on several additional factors that may influence glucose levels in individuals with T1D such as menstrual cycle. Third, although the sample size is large compared with that of other studies with analysis of microbiome composition in individuals with T1D, it may still be insufficient for robust characterization of the bacterial composition and for associating it with PPGR. Finally, we could not evaluate the impact of hybrid closed loop systems on our models, as these systems were not part of the health basket provided by the Israeli Ministry of Health and were not used routinely by individuals with T1D in Israel during the study period.

In summary, here we developed a model that can predict glycemic responses to meals in individuals with T1D, substantially outperforming standard of care. Our work has several potential clinical applications. Our model enables a more accurate prediction of the glycemic response to meals and therefore may allow a better adjustment of the required insulin dosage for meals. It can be further implemented in closed loop systems, personalized decision systems, and alarm systems for the expected high and low blood glucose events for individuals with T1D. These applications have the potential to improve glycemic control, thereby delaying the onset of microvascular complications, while decreasing disease burden. In addition, our model may lead to rationally designed nutritional interventions personally tailored for individuals with T1D based on meals with expected low glycemic response. In addition, our study unravels potential microbial effectors of glycemic control in T1D and underscores the need for additional mechanistic studies that may identify the role of these bacteria and pave the way to novel therapeutic strategies. All of the above have the potential to improve glycemic control and disease management in individuals with T1D.

Acknowledgments. The authors thank the Segal group members for fruitful discussions. Funding. This work is supported by The Israel Science Foundation (ISF) (grant 3-14762). E.S. from the Weizmann institute of Science, Israel is supported by the Crown Human Genome Center, Larson Charitable Foundation New Scientist Fund, Else Kroener Fresenius Foundation, White Rose International Foundation, Ben B. and Joyce E. Eisenberg Foundation, Nissenbaum Family, Marcos Pinheiro de Andrade and Vanessa Buchheim, Lady Michelle Michels, Aliza Moussaieff, and grants funded by the Minerva Foundation with funding from the Federal German Ministry for Education and Research and by the European Research Council and the Israel Science Foundation.

These funding sources had no role in the design of this study or during its execution, analyses, interpretation of the data, or decision to submit results.

**Duality of Interest.** E.S. is a paid consultant for DayTwo. No other potential conflicts of interest relevant to this article were reported.

No pharmaceutical manufacturers or companies from the industry contributed to the planning, design, or conduct of the trial.

Author Contributions. S.S. conceived the project, designed and conducted the analyses, interpreted the results and wrote the manuscript. A.G. designed and conducted the analyses, interpreted the results, and wrote the manuscript. M.R. provided data and interpreted the results. T.Ko. conceived the project, designed the analysis, and interpreted the results. D.K. and T.Ka. designed and conducted the analyses. M.C., N.Z.L., N.S., N.G., N.L., and S.K. provided data and interpreted the results. N.B. designed the analysis and interpreted the results. B.C.W. and Y.G.-G. coordinated and designed data collection. A.W. conceived the project and directed sample sequencing. O.P.-H. and E.S. conceived the project, designed and conducted the analyses, interpreted the results, and supervised the project and analyses. All authors reviewed and approved the manuscript and youch for the accuracy and completeness of the data. E.S. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

#### References

1. Sora ND, Shashpal F, Bond EA, Jenkins AJ. Insulin pumps: review of technological advancement in diabetes management. Am J Med Sci 2019;358:326–331

2. Miller KM, Foster NC, Beck RW, et al.; T1D Exchange Clinic Network. Current state of type 1 diabetes treatment in the U.S.: updated data from the T1D Exchange clinic registry. Diabetes Care 2015;38:971–978

3. Wood JR, Miller KM, Maahs DM, et al.; T1D Exchange Clinic Network. Most youth with type 1 diabetes in the T1D Exchange Clinic Registry do not meet American Diabetes Association or International Society for Pediatric and Adolescent Diabetes clinical guidelines. Diabetes Care 2013;36:2035–2037

4. Aronson R, Brown RE, Abitbol A, et al. The Canadian LMC Diabetes Registry: a profile of the demographics, management, and outcomes of

individuals with type 1 diabetes. Diabetes Technol Ther 2021;23:31–40

5. Nathan DM, Genuth S, Lachin J, et al.; Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. N Engl J Med 1993;329:977–986

6. American Diabetes Association. 9. Pharmacologic approaches to glycemic treatment: *Standards of Care in Diabetes*—2019. Diabetes Care 2019;42(Suppl. 1):S90–S102

 Chase HP, Saib SZ, MacKenzie T, Hansen MM, Garg SK. Post-prandial glucose excursions following four methods of bolus insulin administration in subjects with type 1 diabetes. Diabet Med 2002;19:317–321

8. Bell KJ, Toschi E, Steil GM, Wolpert HA. Optimized mealtime insulin dosing for fat and protein in type 1 diabetes: application of a model-based approach to derive insulin doses for open-loop diabetes management. Diabetes Care 2016;39:1631–1634

9. Wolpert HA, Atakov-Castillo A, Smith SA, Steil GM. Dietary fat acutely increases glucose concentrations and insulin requirements in patients with type 1 diabetes: implications for carbohydrate-based bolus dose calculation and intensive diabetes management. Diabetes Care 2013;36:810–816

10. Zeevi D, Korem T, Zmora N, et al. Personalized nutrition by prediction of glycemic responses. Cell 2015;163:1079–1094

11. Stewart CJ, Ajami NJ, O'Brien JL, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. Nature 2018;562:583–588

12. American Diabetes Association. Standards of medical care in diabetes. In Clinical Practice Recommendations, 2013. Diabetes Care 2013;36 (Suppl. 1):S11–S66

13. Wolever TM, Jenkins DJ. The use of the glycemic index in predicting the blood glucose response to mixed meals. Am J Clin Nutr 1986;43:167–172

14. Vrolix R, Mensink RP. Variability of the glycemic response to single food products in healthy subjects. Contemp Clin Trials 2010;31:5–11 15. Lundberg S, Lee S-I. A unified approach to interpreting model predictions. 22 May 2017 [preprint]. arXiv:1705.07874

16. Lundberg SM, Erion G, Chen H, et al. From local explanations to global understanding with explainable AI for trees. Nat Mach Intell 2020;2: 56–67

17. Lundberg SM, Erion GG, Lee S-I. Consistent individualized feature attribution for tree ensembles. 12 February 2018 [preprint]. arXiv: 1802.03888

18. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods 2012;9: 357–359

19. Pasolli E, Asnicar F, Manara S, et al. Extensive unexplored human microbiome diversity revealed by over 150,000 genomes from metagenomes spanning age, geography, and lifestyle. Cell 2019;176:649–662.e20

20. Korem T, Zeevi D, Suez J, et al. Growth dynamics of gut microbiota in health and disease inferred from single metagenomic samples. Science 2015;349:1101–1106

21. Madsbad S. Impact of postprandial glucose control on diabetes-related complications: how is the evidence evolving? J Diabetes Complications 2016;30:374–385

22. Beck RW, Bergenstal RM, Riddlesworth TD, et al. Validation of time in range as an outcome measure for diabetes clinical trials. Diabetes Care 2019;42:400–405

23. Woldaregay AZ, Årsand E, Walderhaug S, et al. Data-driven modeling and prediction of blood glucose dynamics: machine learning

applications in type 1 diabetes. Artif Intell Med 2019;98:109–134

24. Gingras V, Taleb N, Roy-Fleming A, Legault L, Rabasa-Lhoret R. The challenges of achieving postprandial glucose control using closed-loop systems in patients with type 1 diabetes. Diabetes Obes Metab 2018;20:245–256

25. Alcántara-Aragón V, Gonzalez C, Corcoy R, Ubeda J, Chico A. Carbohydrate-to-insulin ratio in a Mediterranean population of type 1 diabetic patients on continuous subcutaneous insulin infusion therapy. J Diabetes Sci Technol 2015;9:588–592

26. Kuroda A, Yasuda T, Takahara M, et al. Carbohydrate-to-insulin ratio is estimated from 300-400 divided by total daily insulin dose in type 1 diabetes patients who use the insulin pump. Diabetes Technol Ther 2012;14: 1077–1080

27. Turton JL, Raab R, Rooney KB. Lowcarbohydrate diets for type 1 diabetes mellitus: a systematic review. PLoS One 2018;13:e0194987