


מכון ויצמן למדע

WEIZMANN INSTITUTE OF SCIENCE



Host succinate is an activation signal for *Salmonella* virulence during intracellular infection

Document Version:

Accepted author manuscript (peer-reviewed)

Citation for published version:

Rosenberg, G, Yehezkel, D, Hoffman, D, Ciolli Mattioli, C, Fremder, M, Ben-Arosh, H, Vainman, L, Nissani, N, Hen-Avivi, S, Brenner, S, Itkin, M, Malitsky, S, Ohana, E, Ben-Moshe, NB & Avraham, R 2021, 'Host succinate is an activation signal for *Salmonella* virulence during intracellular infection', *Science*, vol. 371, no. 6527, pp. 400-405. <https://doi.org/10.1126/science.aba8026>

Total number of authors:

15

Digital Object Identifier (DOI):

[10.1126/science.aba8026](https://doi.org/10.1126/science.aba8026)

Published In:

Science

License:

Other

General rights

@ 2020 This manuscript version is made available under the above license via The Weizmann Institute of Science Open Access Collection is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognize and abide by the legal requirements associated with these rights.

How does open access to this work benefit you?

Let us know @ library@weizmann.ac.il

Take down policy

The Weizmann Institute of Science has made every reasonable effort to ensure that Weizmann Institute of Science content complies with copyright restrictions. If you believe that the public display of this file breaches copyright please contact library@weizmann.ac.il providing details, and we will remove access to the work immediately and investigate your claim.

1 **Host succinate is a signal for activation of *Salmonella* virulence during**
2 **intracellular infection**

3

4 **Gili Rosenberg¹, Dror Yehezkel¹, Dotan Hoffman¹, Camilla Ciolli Mattioli¹, Moran**
5 **Fremder³, Hadar Ben-Arosh¹, Leia Vainman¹, Noa Nissani¹, Shelly Hen-Avivi¹, Shirley**
6 **Brenner¹, Maxim Itkin², Sergey Malitsky², Ehud Ohana³, Noa Bossel Ben-Moshe¹, Roi**
7 **Avraham^{1,*}**

8 ¹Department of Biological Regulation, Weizmann Institute of Science, 7610001 Rehovot,
9 Israel

10 ²Life Sciences Core Facilities, Weizmann Institute of Science, 7610001 Rehovot, Israel

11 ³Department of Clinical Biochemistry and Pharmacology, Faculty of Health Sciences, Ben-
12 Gurion University of the Negev, 84105, Beer-Sheva, Israel

13

14

15 *Correspondence: roi.avraham@weizmann.ac.il

16

17 **Abstract**

18 Key to the success of intracellular pathogens is the ability to sense and respond to a changing
19 host cell environment. Macrophages exposed to microbial products undergo metabolic
20 changes that drive inflammatory responses. However, the role of macrophage metabolic
21 reprogramming in bacterial adaptation to the intracellular environment has not been explored.
22 Here, using metabolic profiling and dual RNA-seq, we show that succinate accumulation in
23 macrophages is sensed by intracellular *Salmonella* Typhimurium (*S.Tm*) to promote
24 antimicrobial resistance and type III secretion. *S.Tm* lacking the succinate uptake transporter
25 DcuB display impaired survival in macrophages and in mice. Thus, *S.Tm* co-opts the
26 metabolic reprogramming of infected macrophages as a signal that induces its own virulence
27 and survival, providing an additional perspective on metabolic host–pathogen crosstalk.

28

29 **One-sentence summary:** Host metabolic reprogramming, necessary for immune activation,
30 is co-opted by *Salmonella* to induce virulence during intracellular infection

31 Studies of the interactions between *Salmonella enterica* serovar Typhimurium (*S.Tm*) and
32 macrophages have elucidated both a complex virulence program executed by the pathogen
33 and the opposing ability of macrophages to recognize and kill invading bacteria (1). *S.Tm* has
34 evolved to sense the transition from extracellular to intravacuolar environments,
35 characterized by physiological host processes which aim to eliminate the invading pathogen
36 (2). These intracellular cues provide signals that activate bacterial two-component systems,
37 such as PhoPQ and PmrAB, which promote resistance to host antimicrobial peptides (3), and
38 type III secretion systems, located on the *Salmonella* pathogenicity island 2 (SPI-2), which
39 enables intracellular replication (4). Macrophage detection of invading bacteria via pattern
40 recognition receptors results in disruption of the citric acid (TCA) cycle, which shifts host
41 metabolism from oxidative phosphorylation toward aerobic glycolysis with consequent
42 accumulation of several metabolic intermediates (5). Although it is increasingly understood
43 that these metabolic changes are key to the regulation of host proinflammatory responses
44 during infection (6), the impact of host metabolic reprogramming on intracellular
45 pathogens is largely unknown.

46

47 To determine whether host-derived metabolites modulate the intracellular lifestyle of *S.Tm*,
48 we sought to test their effects on *S.Tm* virulence and survival in macrophages. We utilized 2-
49 deoxyglucose (2DG) to inhibit the shift from oxidative phosphorylation (OXPHOS) to
50 glycolysis (5) in activated bone marrow–derived macrophages (BMDMs) infected with *S.Tm*
51 (fig. S1A). We optimized an effective 2DG dose at a low concentration of 1 mM to minimize
52 off-target effects (Fig. 1A and fig. S1, B and C). Additionally, 2DG may lead to a deficient
53 glucose homeostasis of the bacteria that induces a stress response (7). 2DG treatment of *S.Tm*
54 in culture media caused an impaired growth and induction of stress response genes, that was
55 abolished in an *S.Tm* mutant lacking the transporter of 2DG, *ManZ* (8) (fig. S1, D to F and
56 table S1). Thus, 2DG and the $\Delta manZ$ mutant can serve as a tractable model to study the
57 impact of macrophage metabolic reprogramming on intracellular *S.Tm*.

58

59 To assess the interplay of host metabolism and pathogen dynamics in our model, we applied a
60 dual RNA-seq method (9). We curated a list of 977 macrophage genes that were induced in
61 infected cells (table S2). Host responses to infection with the WT or $\Delta manZ$ were similar (fig.
62 S2A) and 2DG did not induce transcriptional changes in uninfected (naïve) cells (fig. S2B).
63 We classified host genes that are upregulated during infection into three classes, based on
64 their expression changes by 2DG (Fig. 1B, fig. S2C and table S2). Importantly, class 2

65 composed of genes downregulated by 2DG, including genes that are differentially expressed
66 by metabolic reprogramming, verified glycolysis inhibition by 2DG (Fig. 1C and fig. S2D).

67

68 To test the impact of inhibiting host OXPHOS-to-glycolysis shift on intracellular *S.Tm*, we
69 curated 960 bacterial genes that were upregulated during infection and differentially
70 expressed with 2DG treatment (fig. S3A and table S3). Expression of stress response genes in
71 WT 2DG-treated cells (fig. S3B), and of *ripA*, part of the itaconic acid degradation operon
72 (*10*) in $\Delta manZ$ strain (fig. S3C), suggested that the macrophage metabolic state is sensed by
73 intracellular *S.Tm*. Importantly, global regulon analysis (table S4) indicated decreased
74 expression in $\Delta manZ$ -infected, 2DG-treated cells of regulons that mediate intracellular
75 survival in macrophages, most prominently the SPI-2 regulon (hypergeometric test, $P < 0.007$,
76 $Q < 0.12$; Fig. 1, D and E and fig. S3, D and E). Thus, there is a clear correlation between the
77 metabolic state of infected macrophages and *S.Tm* virulence gene expression.

78

79 We next analyzed intracellular accumulation of host metabolites during infection with *S.Tm*.
80 Several metabolites, including primary metabolites of the TCA, were significantly changed
81 upon *S.Tm* infection and also reduced by 2DG treatment (Fig. 2A, fig. S4A, and table S5). To
82 test whether *S.Tm* responds to the accumulation of these host metabolites, we performed a
83 virulence assay utilizing the antimicrobial peptide polymyxin B (PMB). Survival of a $\Delta phoP$
84 strain, that is sensitive to PMB challenge (*11*), was significantly restored only in the presence
85 of succinate and itaconate (Fig. 2B and fig. S4B). *S.Tm* grew normally in medium
86 supplemented with succinate, but not with itaconate, at concentrations found within activated
87 macrophages (*12*) (fig. S4, C and D). Thus, we hypothesized that succinate serves as a signal
88 to induce *S.Tm* virulence.

89

90 Using ^{14}C -labeled succinate, we found that *S.Tm* took up and accumulated exogenous
91 succinate (Fig. 2C). We selected a subset of SPI-2 candidate genes whose expression
92 correlated with metabolic remodeling of infected macrophages (from Fig. 1D). We measured
93 induced mRNA expression and fluorescence from transcriptional reporters of these genes in
94 succinate-containing media (Fig. 2, D and E and fig. S4E). Succinate also induced protein
95 levels of one of these SPI-2 effectors, SseL (Fig. 2F). Finally, *S.Tm* grown in succinate-
96 containing media showed increased survival in BMDMs compared to control (Fig. 2G). Thus,
97 succinate is sufficient to mount a bacterial pathogenic program.

98

99 Using RNA-seq, we comprehensively analyzed the direct effects of succinate on *S.Tm* (table
100 S6). Succinate induced the expression of several regulons known to mediate *S.Tm* virulence
101 within macrophages (Fig. 3A, fig. S5A, and table S7). In line with the observed effect of host
102 metabolic shift on induction of SPI-2 genes during intracellular infection, growth in
103 succinate-containing media significantly induced expression of the SPI-2 regulon
104 (hypergeometric test, $Q < 0.001$). Furthermore, succinate induced the expression of the
105 PmrAB regulon (hypergeometric test, $Q < 0.001$), important for antimicrobial peptide
106 resistance (3) (Fig. 3B). We tested whether succinate-induced PMB resistance (Fig. 2B) is
107 mediated by PmrA-regulated genes (Fig. 3C, fig. S5B). Similar to the $\Delta phoP$ strain, the
108 $\Delta pmrA$ strain was susceptible to PMB in control media. However, succinate exposure did not
109 induce PMB resistance in $\Delta pmrA$. Resistance was preserved in a $\Delta phoP$ PmrA^c strain, but not
110 in a $\Delta pmrA$ PhoP^C strain. Thus, succinate-mediated PMB resistance appears to be through
111 induction of PmrA-regulated virulence genes, which deviates from the canonical activation
112 by PhoPQ (13). Succinate also induced resistance to the intracellular antimicrobial peptide,
113 defensin (14) (Fig. 3D and fig. S5C), suggesting a role for succinate in protection against
114 antimicrobial peptides during intracellular infection within macrophages.

115

116 During gut colonization, *S.Tm* can utilize succinate generated by the gut microbiota to
117 complete the TCA cycle (15). To test this possibility, we generated an *S.Tm* double mutant
118 lacking *phoP* and the succinate dehydrogenase complex, subunit A ($\Delta sdhA \Delta phoP$). Similar
119 to $\Delta phoP$, $\Delta sdhA \Delta phoP$ bacteria grown with succinate showed restored resistance to PMB
120 (Fig. 3E and fig. S5, D and E). This result is consistent with succinate acting as a signal for
121 *S.Tm* virulence independent of its role in central metabolism.

122

123 Shifts in host succinate levels might be sensed in the bacterial outer membrane or accumulate
124 within the bacterial cytoplasm. Preferred sugars, such as glycerol, induce carbon catabolite
125 repression, which prevents succinate uptake (16). Bacteria grown with succinate alone
126 exhibited a dose-dependent induction of the SPI-2 gene *steC*, which was lost in media
127 containing glycerol (Fig. 4A and fig. S6, A and B). The addition of permeable succinate
128 (diethylsuccinate) resulted in elevated *steC* expression even in the presence of glycerol, or of
129 media from activated macrophages (fig. S6C). Thus, succinate is actively transported into
130 bacteria in order to induce *S.Tm* pathogenicity.

131

132 Succinate transport into bacteria is mediated by C4-dicarboxylate transporters (17). Growth
133 in succinate-containing media increased expression of three C4-dicarboxylate transporters
134 (Fig. 4B). We examined intracellular survival of the three transporter mutants and measured a
135 significantly lower survival of the $\Delta dcuB$ mutant (Fig. 4C and fig. S6D). Expression of *dcuB*
136 was induced by growth in succinate-containing media and within BMDMs (fig. S6, E and F).
137 Since genetic redundancy may mask individual transporter phenotypes (18), we assessed
138 succinate uptake in a triple *dcu* mutant ($\Delta 3$) background. Succinate uptake was inhibited in
139 the $\Delta 3$ mutant and was significantly restored by complementation of DcuB (Fig. 4D). Similar
140 effects were measured for the expression of the SPI-2 effector *steC* (fig. S6G). In
141 macrophages infected with a WT or $\Delta dcuB$ strains, secretion of SseL SPI-2 effector to the
142 host cytosol was significantly reduced in the $\Delta dcuB$ mutant (Fig. 4E). Thus, active transport
143 of succinate through DcuB is necessary for *S.Tm* virulence within macrophages.

144

145 To test the role of succinate uptake during in vivo infection, mice were intraperitoneally (i.p.)
146 infected either with WT and $\Delta dcuB$ mutant or with a $\Delta phoP$ and a $\Delta phoP \Delta dcuB$ mutants
147 (Fig. 4F). There was no significant difference in the competitive index between the WT and
148 $\Delta dcuB$. By contrast, the $\Delta phoP \Delta dcuB$ had significantly lower survival compared to the
149 $\Delta phoP$ mutant at 48 and 96 hours after infection (Paired *t* test $P < 0.012$, $P < 0.0003$
150 respectively). Thus, the relationship between host accumulation of succinate and bacterial
151 virulence, mediated by DcuB, is also relevant for in vivo infection.

152

153 This work indicates that *S.Tm* has co-opted the metabolic reprogramming of infected
154 macrophages as a signal that induces its own virulence (fig. S7). Other studies reinforce the
155 importance of host and microbiota metabolic products that promote colonization by *S.Tm* and
156 other pathogens (19–21). Our study indicates that during intracellular infection, uptake of
157 host succinate by the DcuB transporter converges on a unique pathway for virulence
158 induction. One plausible mechanism could be by direct interaction of metabolites with
159 bacterial proteins, as with host proteins (5). We suggest a model whereby bacterial virulence
160 is regulated by sensing of TCA intermediates following macrophage metabolic shift, that may
161 be a general theme required for the pathogenic success of diverse intracellular bacteria.

162

163

164 **References and Notes**

- 165 1. E. Mills, R. Avraham, Breaking the population barrier by single cell analysis: one host against
166 one pathogen. *Curr. Opin. Microbiol.* **36**, 69–75 (2017).
- 167 2. D. Drecktrah, L. A. Knodler, R. Ireland, O. Steele-Mortimer, The Mechanism of Salmonella
168 Entry Determines the Vacuolar Environment and Intracellular Gene Expression: The
169 Mechanism of Salmonella Internalization Determines Virulence Gene Expression. *Traffic.* **7**,
170 39–51 (2006).
- 171 3. J. S. Gunn, The Salmonella PmrAB regulon: lipopolysaccharide modifications, antimicrobial
172 peptide resistance and more. *Trends Microbiol.* **16**, 284–290 (2008).
- 173 4. E. Jennings, T. L. M. Thurston, D. W. Holden, Salmonella SPI-2 Type III Secretion System
174 Effectors: Molecular Mechanisms And Physiological Consequences. *Cell Host Microbe.* **22**,
175 217–231 (2017).
- 176 5. D. G. Ryan, L. A. J. O’Neill, Krebs Cycle Reborn in Macrophage Immunometabolism. *Annu.*
177 *Rev. Immunol.* **38**, 289–313 (2020).
- 178 6. L. E. Gleeson, F. J. Sheedy, E. M. Palsson-McDermott, D. Triglia, S. M. O’Leary, M. P.
179 O’Sullivan, L. A. J. O’Neill, J. Keane, Cutting Edge: *Mycobacterium tuberculosis* Induces
180 Aerobic Glycolysis in Human Alveolar Macrophages That Is Required for Control of
181 Intracellular Bacillary Replication. *J. Immunol.* **196**, 2444–2449 (2016).
- 182 7. M. Bobrovskyy, C. K. Vanderpool, The small RNA SgrS: roles in metabolism and pathogenesis
183 of enteric bacteria. *Front. Cell. Infect. Microbiol.* **4** (2014), doi:10.3389/fcimb.2014.00061.
- 184 8. B. Erni, B. Zanolari, H. P. Kocher, The mannose permease of Escherichia coli consists of three
185 different proteins. Amino acid sequence and function in sugar transport, sugar phosphorylation,
186 and penetration of phage lambda DNA. *J. Biol. Chem.* **262**, 5238–5247 (1987).
- 187 9. R. Avraham, N. Haseley, A. Fan, Z. Bloom-Ackermann, J. Livny, D. T. Hung, A highly
188 multiplexed and sensitive RNA-seq protocol for simultaneous analysis of host and pathogen
189 transcriptomes. *Nat. Protoc.* **11**, 1477 (2016).
- 190 10. S. Srikumar, C. Kröger, M. Hébrard, A. Colgan, S. V. Owen, S. K. Sivasankaran, A. D. S.
191 Cameron, K. Hokamp, J. C. D. Hinton, RNA-seq Brings New Insights to the Intra-Macrophage
192 Transcriptome of Salmonella Typhimurium. *PLOS Pathog.* **11**, e1005262 (2015).
- 193 11. Y. Shi, M. J. Cromie, F.-F. Hsu, J. Turk, E. A. Groisman, PhoP-regulated Salmonella resistance
194 to the antimicrobial peptides magainin 2 and polymyxin B. *Mol. Microbiol.* **53**, 229–241 (2004).
- 195 12. L. Tretter, A. Patocs, C. Chinopoulos, Succinate, an intermediate in metabolism, signal
196 transduction, ROS, hypoxia, and tumorigenesis. *Biochim. Biophys. Acta BBA - Bioenerg.* **1857**,
197 1086–1101 (2016).
- 198 13. L. F. F. Kox, M. M. S. M. Wösten, E. A. Groisman, A small protein that mediates the activation
199 of a two-component system by another two-component system. *EMBO J.* **19**, 1861–1872
200 (2000).
- 201 14. P. I. Fields, E. A. Groisman, F. Heffron, A Salmonella locus that controls resistance to
202 microbicidal proteins from phagocytic cells. *Science.* **243**, 1059–1062 (1989).
- 203 15. L. Spiga, M. G. Winter, T. F. de Carvalho, W. Zhu, E. R. Hughes, C. C. Gillis, C. L. Behrendt,
204 J. Kim, D. Chessa, H. L. Andrews-Polymeris, D. P. Beiting, R. L. Santos, L. V. Hooper, S. E.

- 205 Winter, An Oxidative Central Metabolism Enables Salmonella to Utilize Microbiota-Derived
206 Succinate. *Cell Host Microbe*. **22**, 291-301.e6 (2017).
- 207 16. J. L. Botsford, R. D. DeMoss, Catabolite Repression of Tryptophanase in Escherichia coli. *J.*
208 *Bacteriol.* **105**, 303–312 (1971).
- 209 17. I. G. Janausch, E. Zientz, Q. H. Tran, A. Kröger, G. Unden, C4-dicarboxylate carriers and
210 sensors in bacteria. *Biochim. Biophys. Acta BBA - Bioenerg.* **1553**, 39–56 (2002).
- 211 18. A. Kleefeld, B. Ackermann, J. Bauer, J. Krämer, G. Unden, The Fumarate/Succinate Antiporter
212 DcuB of Escherichia coli Is a Bifunctional Protein with Sites for Regulation of DcuS-dependent
213 Gene Expression. *J. Biol. Chem.* **284**, 265–275 (2009).
- 214 19. P. Thiennimitr, S. E. Winter, M. G. Winter, M. N. Xavier, V. Tolstikov, D. L. Huseby, T.
215 Sterzenbach, R. M. Tsois, J. R. Roth, A. J. Bäuml, Intestinal inflammation allows Salmonella
216 to use ethanolamine to compete with the microbiota. *Proc. Natl. Acad. Sci.* **108**, 17480–17485
217 (2011).
- 218 20. A. Jacobson, L. Lam, M. Rajendran, F. Tamburini, J. Honeycutt, T. Pham, W. Van Treuren, K.
219 Pruss, S. R. Stabler, K. Lugo, D. M. Bouley, J. G. Vilches-Moure, M. Smith, J. L. Sonnenburg,
220 A. S. Bhatt, K. C. Huang, D. Monack, A Gut Commensal-Produced Metabolite Mediates
221 Colonization Resistance to Salmonella Infection. *Cell Host Microbe*. **24**, 296-307.e7 (2018).
- 222 21. M. M. Curtis, Z. Hu, C. Klimko, S. Narayanan, R. Deberardinis, V. Sperandio, The Gut
223 Commensal Bacteroides thetaiotaomicron Exacerbates Enteric Infection through Modification
224 of the Metabolic Landscape. *Cell Host Microbe*. **16**, 759–769 (2014).
- 225
- 226 22. A. Rytönen, J. Poh, J. Garmendia, C. Boyle, A. Thompson, M. Liu, P. Freemont, J. C. D.
227 Hinton, D. W. Holden, SseL, a Salmonella deubiquitinase required for macrophage killing and
228 virulence. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 3502–3507 (2007).
- 229
- 230 23. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in Escherichia coli
231 K-12 using PCR products. *Proc. Natl. Acad. Sci.* **97**, 6640–6645 (2000).
- 232
- 233 24. S. Porwollik, C. A. Santiviago, P. Cheng, F. Long, P. Desai, J. Fredlund, S. Srikumar, C. A.
234 Silva, W. Chu, X. Chen, R. Canals, M. M. Reynolds, L. Bogomolnaya, C. Shields, P. Cui, J.
235 Guo, Y. Zheng, T. Endicott-Yazdani, H.-J. Yang, A. Maple, Y. Ragoza, C. J. Blondel, C.
236 Valenzuela, H. Andrews-Polymenis, M. McClelland, Defined Single-Gene and Multi-Gene
237 Deletion Mutant Collections in Salmonella enterica sv Typhimurium. *PLOS ONE*. **9**, e99820
238 (2014).
- 239
- 240 25. S. I. Miller, J. J. Mekalanos, Constitutive expression of the phoP regulon attenuates Salmonella
241 virulence and survival within macrophages. *J. Bacteriol.* **172**, 2485–2490 (1990).
- 242
- 243 26. O. Gal-Mor, D. L. Gibson, D. Baluta, B. A. Vallance, B. B. Finlay, A Novel Secretion Pathway
244 of Salmonella enterica Acts as an Antivirulence Modulator during Salmonellosis. *PLOS*
245 *Pathog.* **4**, e1000036 (2008).
- 246
- 247 27. F. van den Ent, J. Löwe, RF cloning: A restriction-free method for inserting target genes into
248 plasmids. *J. Biochem. Biophys. Methods*. **67**, 67–74 (2006).
- 249
- 250 28. B. K. Coombes, N. F. Brown, Y. Valdez, J. H. Brumell, B. B. Finlay, Expression and Secretion
251 of Salmonella Pathogenicity Island-2 Virulence Genes in Response to Acidification Exhibit

- 252 Differential Requirements of a Functional Type III Secretion Apparatus and SsaL. *J. Biol.*
253 *Chem.* **279**, 49804–49815 (2004).
254
- 255 29. E. L. Mills, B. Kelly, A. Logan, A. S. H. Costa, M. Varma, C. E. Bryant, P. Turlomousis, J. H.
256 M. Däbritz, E. Gottlieb, I. Latorre, S. C. Corr, G. McManus, D. Ryan, H. T. Jacobs, M. Szibor,
257 R. J. Xavier, T. Braun, C. Frezza, M. P. Murphy, L. A. O’Neill, Succinate Dehydrogenase
258 Supports Metabolic Repurposing of Mitochondria to Drive Inflammatory Macrophages. *Cell*.
259 **167**, 457–470.e13 (2016).
260
- 261 30. L. A. Falk, M. M. Hogan, S. N. Vogel, Bone marrow progenitors cultured in the presence of
262 granulocyte-macrophage colony-stimulating factor versus macrophage colony-stimulating
263 factor differentiate into macrophages with distinct tumoricidal capacities. *J. Leukoc. Biol.* **43**,
264 471–476 (1988).
265
- 266 31. S. P. Hmiel, M. D. Snavely, C. G. Miller, M. E. Maguire, Magnesium transport in *Salmonella*
267 typhimurium: characterization of magnesium influx and cloning of a transport gene. *J.*
268 *Bacteriol.* **168**, 1444–1450 (1986).
269
- 270 32. F. Wang, S. Zhang, I. Vuckovic, R. Jeon, A. Lerman, C. D. Folmes, P. P. Dzeja, J. Herrmann,
271 Glycolytic Stimulation Is Not a Requirement for M2 Macrophage Differentiation. *Cell Metab.*
272 **28**, 463–475.e4 (2018).
273
- 274 33. C. Kröger, A. Colgan, S. Srikumar, K. Händler, S. K. Sivasankaran, D. L. Hammarlöf, R.
275 Canals, J. E. Grissom, T. Conway, K. Hokamp, J. C. D. Hinton, An Infection-Relevant
276 Transcriptomic Compendium for *Salmonella enterica* Serovar Typhimurium. *Cell Host*
277 *Microbe.* **14**, 683–695 (2013).
278
- 279 34. A. J. Westermann, K. U. Förstner, F. Amman, L. Barquist, Y. Chao, L. N. Schulte, L. Müller,
280 R. Reinhardt, P. F. Stadler, J. Vogel, Dual RNA-seq unveils noncoding RNA functions in host–
281 pathogen interactions. *Nature.* **529**, 496–501 (2016).
282
- 283 35. S. Malitsky, C. Ziv, S. Rosenwasser, S. Zheng, D. Schatz, Z. Porat, S. Ben-Dor, A. Aharoni, A.
284 Vardi, Viral infection of the marine alga *Emiliana huxleyi* triggers lipidome remodeling and
285 induces the production of highly saturated triacylglycerol. *New Phytol.* **210**, 88–96 (2016)
286
- 287 36. M. Salem, M. Bernach, K. Bajdzienko, P. Giavalisco, A Simple Fractionated Extraction
288 Method for the Comprehensive Analysis of Metabolites, Lipids, and Proteins from a Single
289 Sample. *J. Vis. Exp. JoVE* (2017), doi:10.3791/55802.
290
- 291 37. L. Zheng, S. Cardaci, L. Jerby, E. D. MacKenzie, M. Sciacovelli, T. I. Johnson, E. Gaude, A.
292 King, J. D. G. Leach, R. Edrada-Ebel, A. Hedley, N. A. Morrice, G. Kalna, K. Blyth, E.
293 Ruppin, C. Frezza, E. Gottlieb, Fumarate induces redox-dependent senescence by modifying
294 glutathione metabolism. *Nat. Commun.* **6**, 6001 (2015).
295
- 296 38. A. P. R. Bally, P. Lu, Y. Tang, J. W. Austin, C. D. Scharer, R. Ahmed, J. M. Boss, NF-κB
297 Regulates PD-1 Expression in Macrophages. *J. Immunol.* **194**, 4545–4554 (2015).
298
- 299 39. D. G. Russell, L. Huang, B. C. VanderVen, Immunometabolism at the interface between
300 macrophages and pathogens. *Nat. Rev. Immunol.* **19**, 291 (2019).
301
- 302 40. B. Kelly, L. A. J. O’Neill, Metabolic reprogramming in macrophages and dendritic cells in
303 innate immunity. *Cell Res.* **25**, 771–784 (2015).

305 **Acknowledgements**

306 We thank A. Barczak, E. Gottlieb, and A. Vardi for their valuable advice and their reading
307 and commentary on this manuscript. This work was supported by the European Research
308 Committee (ERC grant No. 756653), the Israel Science Foundation (grant No. 1890/17), the
309 Minerva Foundation with funding from the Federal Ministry for Education and Research, the
310 Estate of David Turner, the Merle S. Cahn Foundation, the Estate of Sylvia Holder, the Estate
311 of Zvia Zeroni and the Estate of Leah Arbel. R.A. is the Incumbent of Philip Harris and
312 Gerald Ronson career development chair.

313

314 **Author contributions**

315 G.R. and R.A. designed the study. G.R., D.H., C.C.M., H.B.A., L.V., S.H.A., and S.B.
316 performed the experiments. M.F. and E.O. performed ¹⁴C succinate uptake experiments. M.I.
317 and S.M. performed metabolomics experiments. D.Y. and N.N. analyzed the data. G.R. and
318 R.A. wrote the manuscript.

319

320 **Competing Interests**

321 The authors declare no competing interests

322

323 **Data and materials availability**

324 All RNA-seq data have been deposited in NCBI's Gene Expression Omnibus (GEO) under
325 the super-series accession number GSE139208. All data are available in the manuscript or the
326 supplementary materials.

327

328 **Supplementary Materials**

329 Material and Methods

330 Figures S1-S7

331 Tables S1-S10

332 Supplementary references (22-40)

333

334 **Figure legends**

335

336 **Figure 1: Interplay between macrophage metabolic reprogramming and *S. Tm***
337 **virulence.**

338 (A) BMDMs were infected with *S. Tm* and treated with 2DG; basal respiration (OCR, gray)
339 and stimulated glycolysis (ECAR, black) were measured. Error bars indicate means \pm SEM of
340 3-6 replicates representative of three independent experiments. (B) Heatmap of host genes
341 upregulated in *S. Tm*-infected BMDMs compared to uninfected controls. Genes were
342 classified into three classes based on their expression changes with 2DG. Colorbars: relative
343 gene expression (red–blue) and fold change between \pm 2DG (yellow–purple). (C and D) Host
344 class 2 genes and bacterial SPI-2 genes presented as the calculated difference in expression
345 between 2DG-treated and untreated samples. (E) MA plots of bacterial transcripts in 2DG-
346 treated over untreated samples (y -axis) versus the average absolute read counts (x -axis).

347

348 **Figure 2: *S. Tm* virulence is induced in response to succinate.**

349 (A) BMDMs were infected with a $\Delta manZ$ strain, treated with 2DG, and intracellular levels of
350 central metabolites were measured by mass spectrometry. (B) $\Delta phoP$ mutant grown without
351 (control; gray) or with PMB (blue) supplemented with 10 mM of the indicated metabolites.
352 Survival was measured by CFU. (C) *S. Tm* was cultured in the indicated media supplemented
353 with ^{14}C labeled succinate, and succinate uptake was measured. Background indicates heat-
354 killed bacteria. (D-G) *S. Tm* was grown in control media or media supplemented with 10 mM
355 succinate. The expression of *steC*, *sseL*, and *gapA* (normalizing control) was measured by
356 quantitative real-time PCR (D). Fluorescence signal was measured from *S. Tm* strains
357 expressing a transcriptional reporter fused to selected SPI-2 promoters (normalized to OD₆₀₀)
358 (E). Immunoblot of SseL protein (SseL-2HA) and GroEL (F). BMDMs were infected with
359 *S. Tm* and intracellular survival was measured by CFU (G).
360 Error bars indicate means \pm SEM, representative of two [(C)] or three [(B), (D-G)]
361 independent experiments. Unpaired t test: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

362

363 **Figure 3: Succinate induces SPI-2 and antimicrobial resistance.**

364 (A) RNA-seq analysis of *S. Tm* grown in control media or succinate-containing media. SPI-2
365 and PmrAB regulons were significantly induced in succinate. (B) mRNA expression of genes
366 induced by succinate (taken from A) across the PMB resistance locus. (C to E) *S. Tm* strains
367 were grown in the presence of PMB (C and E) or defensin (D), and survival measured by
368 CFU. Error bars indicate means \pm SEM, representative of three independent experiments.
369 Unpaired t test: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

370

371 **Figure 4: Uptake of host succinate is required for intracellular survival of *S. Tm*.**

372 (A) *S. Tm* harboring a PsteC-*cfp* reporter was grown in media without or with glycerol, and
373 succinate or diethylsuccinate as indicated, and fluorescence was measured (normalized to
374 OD₆₀₀). (B) Expression of the C4-dicarboxylate transporter genes taken from RNA-seq data of
375 Fig. 3A. (C) BMDMs were infected with the indicated *S. Tm* strains and intracellular survival
376 was measured by CFU. (D) *S. Tm* strains were cultured in succinate-containing media, and
377 succinate uptake was measured by ^{14}C labeled succinate. (E) Macrophages were infected
378 with WT or $\Delta dcuB$ strains. SseL protein was quantified by protein immunoblot of bacterial
379 pellet and host cell cytosol fractions. Boxplots: SseL protein abundance. (F) Groups of mice
380 (boxes) were infected i.p. with either WT and $\Delta dcuB$ or $\Delta phoP$ and $\Delta phoP \Delta dcuB$ mutants,
381 and competitive index was measured by CFU.

382 Error bars indicate means \pm SEM, representative of two [(B), (D), (F)] or three [(A), (C), (E)]
383 independent experiments. Unpaired *t* test: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.
384