

מכון ויצמן למדע

WEIZMANN INSTITUTE OF SCIENCE



Are Bone Marrow-Derived Fibroblastoid Colony-Forming Cells Involved in Viral Leukemogenesis?

Document Version:

Publisher's PDF, also known as Version of record

Citation for published version:

Zipori, D 1977, Are Bone Marrow-Derived Fibroblastoid Colony-Forming Cells Involved in Viral Leukemogenesis? in P Bentvelzen, J Hilgers & DS Yohn (eds), *Advances in Comparative Leukemia Research: proceedings of the VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, held in Amsterdam, the Netherlands, on 22-26 August 1977*. Amsterdam, pp. 314-315, VIIIth International Symposium on Comparative Research on Leukemia and Related Diseases, Amsterdam, Netherlands, 22/8/77.

Total number of authors:

1

Published In:

Advances in Comparative Leukemia Research

License:

Other

General rights

@ 2020 This manuscript version is made available under the above license via The Weizmann Institute of Science Open Access Collection is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognize and abide by the legal requirements associated with these rights.

How does open access to this work benefit you?

Let us know @ library@weizmann.ac.il

Take down policy

The Weizmann Institute of Science has made every reasonable effort to ensure that Weizmann Institute of Science content complies with copyright restrictions. If you believe that the public display of this file breaches copyright please contact library@weizmann.ac.il providing details, and we will remove access to the work immediately and investigate your claim.

ARE BONE MARROW DERIVED FIBROBLASTOID COLONY-FORMING CELLS INVOLVED IN VIRAL LEUKEMOGENESIS?

D. Zipori

Radiobiological Institute TNO, Lange Kleiweg 151, Rijswijk (The Netherlands)

Haemopoietic tissues contain cells capable of forming fibroblastoid colonies in vitro. It has been suggested that these cells belong to the stromal component of the haemopoietic tissue and that they induce specific haemopoietic microenvironment¹. We assayed the incidence of bone marrow derived fibroblastoid colony-forming cells in the BALB/c, C3H, C57BL and (BALB/c x DBA/2)F1 (CD2) mouse strains. The number of colony-forming cells declined from 82 ± 19 per 10^6 bone marrow cells at birth, to 27 ± 5 per 10^6 bone marrow cells in 3 week old mice and 19 ± 6 per 10^6 in 6 week old mice (fig. 1). The above mentioned strains are all characterized by low spontaneous leukaemia rates. Inoculation of Soule tumour virus into 5 day old BALB/c mice induced high lymphosarcoma incidence with an average latency of 4 months. As early as 2 weeks following virus inoculation, a reduction in the number of bone marrow derived fibro-




Fig. 1. Incidence of fibroblastoid colony forming cells in mouse bone marrow.

blastoid colony-forming cells was observed (fig. 1). In contrast, the total cellularity and the number of CFU-s and CFU-c in the bone marrow did not differ significantly from the numbers in control animals.

A low incidence of bone marrow derived fibroblastoid colony-forming cells was also observed in AKR mice which are characterized by a high spontaneous leukaemia rate. Using a modified XC plaque assay we found that fibroblastoid colonies from both AKR mice and from Soule virus inoculated BALB/c mice harboured an ecotropic virus. In the latter mouse strain budding C type particles were observed in electron microscope preparations of cells from fibroblastoid colonies. The presence of ecotropic virus in fibroblastoid cells did not seem to be part of a generalized infection of the bone marrow, since

cells in CFU-c colonies assayed under similar conditions had very low XC transforming activity (Table 1).

The results presented here demonstrate a correlation between high leukaemia incidence in mice (AKR and Soule virus infected BALB/c) and ecotropic virus producing fibroblastoid cells in the bone marrow. RFM mice were also found to have a low incidence of fibroblastoid colony forming cells early in life (fig. 1). This mouse strain shows a high incidence of lymphoreticular tumours in old age². However, no evidence was obtained so far for early expression of virus in the bone marrow of this mouse strain.

If fibroblastoid colony-forming cells and their progeny participate in the regulation of haemopoiesis, their infection by leukaemia viruses could be the cause of pre-leukaemic perturbations of haemopoiesis. Moreover, these cells may serve as a site for malignant transformation of haemopoietic cells by simultaneously providing differentiation signals and oncogenic viruses.

TABLE 1.

PRESENCE OF XC SYNCYTIA INDUCING VIRUS IN BONE MARROW DERIVED FIBROBLASTOID COLONIES FROM AKR MICE

mouse strain	number of cells/plate	CSF	number of colonies /plate and colony type	number of XC syncytia/plate*	number of XC syncytia/colony (calculated)
AKR	5×10^4	+	60 ± 10 (CFU-c)	20 ± 10	0.3
	5×10^4	-	0	17 ± 5	-
	6×10^5	-	10 ± 5 (fibroblastoid)	500 ± 50	50.0
BALB/c	5×10^4	+	70 ± 15 (CFU-c)	0	0
	5×10^4	-	0	0	-
	6×10^5	-	35 ± 11 (fibroblastoid)	0	0

* XC cells were plated on top of preformed fibroblastoid or CFU-c colonies and syncytia scored 4 days later.

REFERENCES

1. Friedenstein, A.J., Chailakhjan, R.K., Latsinik, N.V., Panasyuk, A.F. and Keiliss-Borok, I.V. (1974) *Transplantation*, 17, 331-340.
2. Zurcher, C., van Zwieten, M.J. and Hollander, C.F. (1976) In *REP Annual Report*, pp. 280-281.

ACKNOWLEDGEMENT

I am greatly indebted to Mrs. M.J. Molenaar for her invaluable technical assistance. This study has been supported by the Koningin Wilhelmina Fonds, The Netherlands Organization for the Fight against Cancer.